

FIG. 1




FIG. 2




FIG. 3




FIG. 4A




FIG. 4B




FIG. 5A




FIG. 5B




FIG. 6




FIG. 7




FIG. 8




FIG. 9




FIG. 10




FIG. 11




FIG. 12




FIG. 13




FIG. 14




FIG. 15




FIG. 16





FIG. 17



MAGNIFICATION : X 1,000,000 TIMES

66260"02004520

FIG. 18



20nm

20250415 09:25:25

FIG. 19




FIG. 20





FIG. 21



MAGNIFICATION: X 1,000,000 TIMES

Figure 1 consists of 12 bar charts, labeled (a) through (l), each representing a different fish species. The species are: (a) Atlantic croaker, (b) Striped bass, (c) Weakfish, (d) Bluefish, (e) Spot, (f) Rockfish, (g) Atlantic silverside, (h) Atlantic herring, (i) Atlantic menhaden, (j) Atlantic bluefish, (k) Atlantic tomcod, and (l) Atlantic silverside. Each chart shows the percentage of total catch for that species from 1990 to 2001. The y-axis for all charts is 'Percentage of total catch' and ranges from 0 to 100. The x-axis is 'Year' and ranges from 1990 to 2001. Each bar represents the mean percentage for that year, and error bars indicate the standard error. The data shows varying trends for each species over the 12-year period.




FIG. 23





FIG. 24



AVERAGE GRAIN DIAMETER : 10 TO 20 nm

FIG. 25




FIG. 26




FIG. 27




FIG. 28




FIG. 29




FIG. 30




FIG. 31




FIG. 32




FIG. 33




FIG. 34




FIG. 35




FIG. 36




FIG. 37




FIG. 38

